3 (Sem-5/CBCS) PHY HE 3

2024

PHYSICS

(Honours Elective)

Paper: PHY-HE-5036

(Advanced Mathematical Physics-I)

Full Marks: 60

Time: Three hours

The figures in the margin indicate full marks for the questions.

- 1. Answer the following questions: $1 \times 7 = 7$
 - (a) What do you mean by basis of a vector space?
 - (b) How can be obtained an orthonormal set of vectors from an orthogonal set?
 - (c) What is called an abelian group?
 - (d) Find ln A, where $A = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$.
 - (e) What is Einstein's summation

convention?

- (f) Show that $\delta_{ij}\varepsilon_{ijk}=0$.
- (g) If A_i and A' represent first rank covariant and contravariant tensors respectively, prove that $A_i = A'$ in Cartesian coordinate system.
- 2. Answer the following questions: $2\times4=8$
 - (a) Determine whether the vectors (1, 2, 3) and (2,-2,0) are linearly independent or not.
 - (b) State and verify Cayley-Hamilton theorem for the matrix $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$. 1+1=2
 - (c) Using tensor notations, prove that $\vec{\nabla} \times (\vec{\nabla} \phi) = 0$
 - (d) What is Minkowski space? What are the transformation equations relating coordinates in this space? 1+1=2

- 3. Answer **any three** questions from the following: 5×3=15
 - (a) (i) Diffine a group by meationing axioms. $2\frac{1}{2}$
 - (ii) Show that the set of $n \times n$ unitary matrices forms a group under matrix multiplication. $2\frac{1}{2}$
 - (b) (i) Using tensor notations, show that $\operatorname{div} \vec{A}$ is an invariant.
 - (ii) Prove that diagonalizing matrix of a real symmetric matrix is orthogonal. 2
 - (c) (i) Using direction cosines, establish the relation $\overline{x}_i = a_{ii} x_i$.
 - (ii) Write the inverse transformation equation.

4+1=5

(d) (i) State quotient law of tensors. 2

3

(ii) Prove that the sum of two tensors of the same rank and type is also a tensor of same rank and type.

3

- (e) (i) Show that the properly of symmetry of a tensor between a pair of dissimilar indices is not invariant under coordinate transformation.
 - (ii) If A^{ij} is an anti-symmetric tensor and B_i is a vector, show that $A^{ij}B_iB_j=0.$
- 4. Answer the following question: (a) or (b), (c) or (d) and (e) or (f) 10×3=30
 - (a) (i) Show that the set of all complex numbers form a vector space over the field of real numbers.
 - (ii) What is the dimension of above mentioned vector space? Justify your answer. 1+2=3
 - (iii) Show that $A^k = ED^k E^{-1}$, where k is any integer, D and E are diagonal and diagonalizing matrices of matrix A.

(b) (i) Evaluate
$$e^{A}$$
, where $A = \begin{bmatrix} \frac{3}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{3}{2} \end{bmatrix}$

(ii) Find the standard matrix of linear transformation T from R^2 to R^4 such that $T(e_1) = (3,1,3,1), T(e_2) = (-5,2,0,0),$ where $e_1 = (1,0)$ and $e_2 = (0,1)$.

(c) (i) Solve the following coupled differential equations:

$$\frac{dx}{dt} = x + y \text{ and}$$

$$\frac{dy}{dt} = 4x + y$$
using method of matrices where $x(0) = y(0) = 1$.

(ii) Find the anti-symmetric tenser of rank two associated with the vector(x,x+y,x+y+z) in three-dimensional space.

- (d) (i) Establish the relation $dS^2 = g_{ij} dx^i dx^j, \text{ where symbols}$ have their usual meanings. 4
 - (ii) Show that $\varepsilon_{iks} \ \varepsilon_{mps} = \delta_{im} \ \delta_{kp} \delta_{ip} \delta_{km} = 0. \qquad 3$
 - (iii) If A^{λ} $B_{\mu\nu}$ is a tensor for all first rank contravariant tensors A^{λ} then show that $B_{\mu\nu}$ is also a tensor. 3
- (e) (i) Using tensor analysis, prove the following vector identities:

$$\vec{\nabla} \cdot (\vec{A} \times \vec{B}) = \vec{B} \cdot (\vec{\nabla} \times \vec{A}) - \vec{A} \cdot (\vec{\nabla} \times \vec{B})$$
 and

$$\vec{A} \times (\vec{B} \times \vec{C}) = \vec{B} \cdot (\vec{A} \times \vec{C}) - \vec{C} \cdot (\vec{A} \cdot \vec{B})$$
2+3=5

(ii) Using tensor analysis, establish the relation $L_i = \varepsilon_{ijk} I_{jk}$ where symbals have their usual meanings.

(iii) If the length of a vector is invariant under coordinate transformation (rotation), show that $a_{ij}a_{ij} = \delta_{ij}$. 2

Or

- (i) Derive with seat diagrams, the components of stress at a point of a solid body in three-dimensional space.
- (ii) Use tensor analysis to find the components of a vector in plane polar coordinates whose components in Cartesian coordinates are \dot{x} and \dot{y} . 5