

Total number of printed pages-8

3 (Sem-5/CBCS) MAT HC 2

2023

## MATHEMATICS

(Honours Core)

Paper : MAT-HC-5026

(Linear Algebra)

Full Marks : 80

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer the following questions as directed :

1×10=10

(a) Let  $A = \begin{pmatrix} 1 & -3 & -2 \\ -5 & 9 & 1 \end{pmatrix}$  and  $\vec{u} = \begin{bmatrix} 5 \\ 3 \\ -2 \end{bmatrix}$ .

Check whether  $\vec{u}$  is in null space of A.

(b) Define subspace of a vector space.

(c) Give reason why  $\mathbb{R}^2$  is not a subspace of  $\mathbb{R}^3$ .

Contd.

(d) State whether the following statement is true **or** false :

“If dimension of a vector space  $V$  is  $p > 0$  and  $S$  is a linearly dependent subset of  $V$ , then  $S$  contains more than  $p$  elements.”

(e) If  $\vec{x}$  is an eigenvector of  $A$  corresponding to the eigenvalue  $\lambda$  then what is  $A^3 \vec{x}$  ?

(f) When two square matrices  $A$  and  $B$  are said to be similar ?

(g) If  $\vec{v} = (1 \ -2 \ 2 \ 4)$  then find  $\|\vec{v}\|$ .

(h) Find a unit vector in the direction of

$$\vec{u} = \begin{bmatrix} 8/3 \\ 2 \end{bmatrix}.$$

(i) Under what condition two vectors  $\vec{u}$  and  $\vec{v}$  are orthogonal to each other ?

(j) Define orthogonal complement of vectors.

2. Answer the following questions :

2×5=10

(a) Show that the set  $W = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} : xy \geq 0 \right\}$  is not a subspace of  $\mathbb{R}^2$ .

(b) Let  $\vec{b}_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $\vec{b}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$ ,  $\vec{x} = \begin{bmatrix} 4 \\ 5 \end{bmatrix}$  and

$\beta = \{b_1, b_2\}$ . Find the coordinate vector  $[x]_\beta$  of  $\vec{x}$  relative to  $\beta$ .

(c) Find the eigenvalues of  $A = \begin{bmatrix} 2 & 3 \\ 3 & -6 \end{bmatrix}$ .

(d) Let  $P_2$  be the vector space of all polynomials of degree less than equal to 2. Consider the linear transformation

$T : P_2 \rightarrow P_2$  defined by

$T(a_0 + a_1t + a_2t^2) = a_1 + 2a_2t$ . Find the matrix representation  $[T]_\beta$  of  $T$  with respect to the base  $\beta = \{1, t, t^2\}$ .

(e) Show that the matrix  $A = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{2}{3} \\ \frac{1}{\sqrt{2}} & -\frac{2}{3} \\ 0 & \frac{1}{3} \end{bmatrix}$

has orthogonal columns.

3. Answer **any four** questions :  $5 \times 4 = 20$

(a) Let  $S = \{v_1, v_2, \dots, v_p\}$  be a set in the vector space  $V$  and  $H = \text{span}(S)$ . Now if one of the vector in  $S$ , say  $v_k$ , is linear combination of the other vectors in  $S$ , then show that  $S$  is linearly dependent and the subset of  $S_1 = S - \{v_k\}$  still span  $H$ .  $2+3=5$

(b) Show that the set of all eigenvectors corresponding to the distinct eigenvalues of a  $n \times n$  matrix  $A$  is linearly independent.

(c) Let  $W$  be a subspace of the vector space  $V$  and  $S$  is a linearly independent subset of  $W$ . Show that  $S$  can be extended, if necessary, to form a basis for  $W$  and  $\dim W \leq \dim V$ .

(d) If  $A = \begin{bmatrix} 1 & 3 & 3 \\ -3 & -5 & -3 \\ 3 & 3 & 1 \end{bmatrix}$ . Find an

invertible matrix  $P$  and a diagonal matrix  $D$  such that  $A = PDP^{-1}$ .

(e) If  $\bar{y} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$  and  $\bar{u} = \begin{bmatrix} 4 \\ -7 \end{bmatrix}$  then find the orthogonal projection of  $\bar{y}$  onto  $\bar{u}$  and write  $\bar{y}$  as the sum of two orthogonal vectors, one in  $\text{span}\{\bar{u}\}$  and the other orthogonal to  $\bar{u}$ .

(f) If  $W = \text{span}\{x_1, x_2\}$  where  $x_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ ,

$x_2 = \begin{bmatrix} \frac{1}{3} \\ \frac{1}{3} \\ -\frac{2}{3} \end{bmatrix}$ , find a orthogonal basis for

$W$ .

Answer **either (a) or (b)** from each of the following questions :  $10 \times 4 = 40$

4. (a) Find a spanning set for the null space of the matrix :

$$A = \begin{bmatrix} -3 & 6 & -1 & 1 & -7 \\ 1 & -2 & 2 & 3 & -1 \\ 2 & -4 & 5 & 8 & -4 \end{bmatrix}$$

Is this spanning set linearly independent?

$8+2=10$

(b) (i) If a vector space  $V$  has a basis of  $n$  vectors, then show that every basis of  $V$  must consist of exactly  $n$  vectors. 4

(ii) Find a basis for column space of the following matrix : 6

$$B = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}$$

5. (a) Define eigenvalue and eigenvector of a matrix. Find the eigenvalues and corresponding eigenvectors of the

matrix  $\begin{bmatrix} 2 & 3 \\ 3 & -6 \end{bmatrix}$ . 2+8=10

(b) Let  $T$  be a linear operator on a finite dimensional vector space  $V$  and let  $W$  denote the  $T$ -cyclic subspace of  $V$  generated by a non-zero vector  $v \in V$ . If  $\dim(W) = k$  then show that

(i)  $\{v, T(v), T^2(v), \dots, T^{k-1}(v)\}$  is a basis for  $W$ .

(ii) If  $a_0v + a_1T(v) + \dots + a_{k-1}T^{k-1}(v) + T^k(v) = 0$ , then the characteristics polynomial of  $T_w$  is

$$f(t) = (-1)^k (a_0 + a_1t + \dots + a_{k-1}t^{k-1} + t^k).$$

6+4=10

6. (a) (i) Define orthogonal set of vectors.

Let  $S = \{\bar{u}_1, \bar{u}_2, \dots, \bar{u}_p\}$  is an orthogonal set of non-zero vectors in  $\mathbb{R}^n$ , then show that  $S$  is linearly independent. 1+4=5

(ii) For any symmetric matrix show that any two eigenvectors from different eigenspaces are orthogonal. 5

(b) Define inner product space. Show that the following is an inner product in  $\mathbb{R}^2$

$$\langle u, v \rangle = 4u_1v_1 + 5u_2v_2$$

Where  $u = (u_1, u_2), v = (v_1, v_2) \in \mathbb{R}^2$

Also, show that in any inner product space  $V$ ,

$$|\langle u, v \rangle| \leq \|u\| \cdot \|v\|, \quad \forall u, v \in V.$$

2+4+4=10

7. (a) (i) Consider the bases  $\beta = \{b_1, b_2\}$  and  $\gamma = \{c_1, c_2\}$  for  $\mathbb{R}^2$  where

$$b_1 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}, \quad b_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}, \quad c_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix}$$

and  $c_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}$ , find the change

of coordinate matrix from  $\gamma$  to  $\beta$  and from  $\beta$  to  $\gamma$ . 5

(ii) Compute  $A^{10}$  where

$$A = \begin{bmatrix} 4 & -3 \\ 2 & -1 \end{bmatrix}. \quad \text{5}$$

(b) State Cayley-Hamilton theorem for matrices. Verify the theorem for the

matrix  $M = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$  and hence find  $M^{-1}$ .