

Total number of printed pages-23

**3 (Sem-5/CBCS) MAT HE 1/2/3**

**2023**

**MATHEMATICS**

(Honours Elective)

**Answer the Questions from any one Option.**

**OPTION-A**

Paper : MAT-HE-5016

**(Number Theory)**

**OPTION-B**

Paper : MAT-HE-5026

**(Mechanics)**

**OPTION-C**

Paper : MAT-HE-5036

**(Probability and Statistics)**

Full Marks : 80

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

Contd.

### OPTION-A

Paper : MAT-HE-5016

#### ( Number Theory )

1. Answer the following questions as directed:

1×10=10

(a) Which of the following Diophantine equations cannot have integer solutions ?

(i)  $33x + 14y = 115$

(ii)  $14x + 35y = 93$

(b) State whether the following statement is true **or** false :

“If  $a$  and  $b$  are relatively prime positive integers, then the arithmetic progression  $a, a + b, a + 2b, \dots$  contains infinitely many primes.”

(c) For any  $a \in \mathbb{Z}$  prove that  $a \equiv a \pmod{m}$ , where  $m$  is a fixed integer.

(d) Under what condition the  $k$  integers  $a_1, a_2, \dots, a_k$  form a CRS  $\pmod{m}$  ?

(e) Find  $\sigma(p)$  where  $p$  is a prime number.

(f) Define Euler's phi function.

(g) If  $n = 12789$ , find  $\tau(n)$ .

(h) If  $x$  is a real number then show that  $[x] \leq x < [x] + 1$ , where  $[ ]$  represents the greatest integer function.

(i) Calculate the exponent of the highest power of 5 that divides  $1000!$

(j) When an arithmetic function  $f$  is said to be multiplicative ?

2. Answer the following questions : 2×5=10

(a) Show that there is no arithmetic progression  $a, a + b, a + 2b, \dots$  that consists solely of prime numbers.

(b) Use properties of congruence to show that 41 divides  $2^{20} - 1$ .

(c) Let  $p > 1$  be a positive integer having the property that  $p \mid a \cdot b \Rightarrow p \mid a$  or  $p \mid b$ , then prove that  $p$  is a prime.

(d) If  $a$  is a positive integer and  $q$  is its least positive divisor then show that  $q \leq \sqrt{a}$ .

(e) For  $n \geq 3$ , evaluate  $\sum_{k=1}^n \mu(k!)$ , here  $\mu$  is the Mobius function.

3. Answer **any four** questions : 5×4=20

(a) If  $(m, n) = 1$  and  $S_1 = \{x_0, x_1, x_2, \dots, x_{m-1}\}$  is a CRS (mod  $m$ ) and

$S_2 = \{y_0, y_1, y_2, \dots, y_{n-1}\}$  is a CRS (mod  $n$ ) then show that the set

$S = \{nx_i + my_j : 0 \leq i \leq m-1, 0 \leq j \leq n-1\}$  form a CRS (mod  $mn$ ).

(b) Find all integers that satisfy simultaneously

$$x \equiv 5 \pmod{18}; x \equiv -1 \pmod{24};$$

$$x \equiv 17 \pmod{33}$$

(c) If  $n \geq 1$  is an integer then show that  $\sigma(n)$  is odd if and only if  $n$  is a perfect square or twice a perfect square.

(d) If  $a_1, a_2, \dots, a_k$  form a RRS (mod  $m$ ) ie. Reduced Residue System modulo  $m$  then show that  $k = \phi(m)$ .

(e) If  $x$  and  $y$  be real numbers then show that  $[x+y] = [x] + [y]$  and  $[-x-y] = [-x] + [-y]$  if and only if one of  $x$  or  $y$  is an integer.

(f) For  $n > 2$ , show that  $\phi(n)$  is an even integer. Here,  $\phi$  is the Euler phi function.

Answer **either (a) or (b)** from each of the following questions : 10×4=40

4. (a) (i) Show that every positive integer can be expressed as a product of primes. Also show that apart from the order in which prime factors occur in the product, they are unique. 3+4=7

(ii) If  $k$  integers  $a_1, a_2, \dots, a_k$  form a CRS (mod  $m$ ), then show that  $m = k$ . 3

(b) (i) Show that any natural number greater than one must have a prime factor. 5

- (ii) Prove that if all the  $n > 2$  terms of the arithmetic progression  $p, p+d, p+2d, \dots, p+(n-1)d$  are prime numbers, then the common difference  $d$  is divisible by every prime  $q < n$ . 5

5. (a) State and prove Wilson's theorem. Is the converse also true? Justify your answer.

$$1+6+3=10$$

- (b) Let  $a$  and  $m > 0$  be integers such that  $(a, m) = 1$ , then show that  $a^{\phi(m)} \equiv 1 \pmod{m}$ , here  $\phi$  is the Euler's phi function. Deduce from it the Fermat's Little theorem. Also find the last two digits of  $3^{1000}$ .

$$5+2+3=10$$

6. (a) For each positive integer  $n \geq 1$ , show that

$$\phi(n) = \sum_{d/n} \mu(d) \frac{n}{d} = n \prod_{p/n} \left(1 - \frac{1}{p}\right)$$

- (b) (i) If  $f$  and  $g$  are two arithmetic functions, then show that the following conditions (A) and (B) are equivalent 7

$$(A) \quad f(n) = \sum_{d/n} g(d)$$

$$(B) \quad g(n) = \sum_{d/n} \mu(d) f\left(\frac{n}{d}\right) = \sum_{d/n} \mu\left(\frac{n}{d}\right) f(d)$$

- (ii) If  $f$  is a multiplicative arithmetic function, then show that

$$g_1(n) = \sum_{d/n} f(d) \quad \text{and}$$

$$g_2(n) = \sum_{d/n} \mu(d) f(d) \quad \text{are both}$$

multiplicative arithmetic functions. 3

7. (a) State and prove Chinese Remainder theorem. 2+8=10

- (b) (i) For  $n > 1$ , show that the sum of the positive integers less than  $n$  and relatively prime to  $n$  is  $\frac{1}{2}n\phi(n)$ . 5

(ii) If  $n \geq 1$  is an integer then show that

$$\prod_{d/n} d = n^{\frac{\tau(n)}{2}}. \text{ Is } \prod_{d/n} d \text{ an integer}$$

when  $\tau(n)$  is odd? Justify. 5

### OPTION-B

Paper : MAT-HE-5026

#### (Mechanics)

1. Answer the following questions :  $1 \times 10 = 10$
- (a) What is the resultant of two equal forces acting at an angle  $120^\circ$ ?
  - (b) State Lami's theorem.
  - (c) State the principle of conservation of linear momentum.
  - (d) When two parallel forces cannot be compounded into a single resultant force?
  - (e) Define impulsive force with an example.
  - (f) State a necessary and sufficient condition for a system of coplanar forces acting on a rigid body to maintain equilibrium.
  - (g) Define amplitude and frequency of a simple harmonic motion (SHM).
  - (h) Write down the relation between the angle of friction and co-efficient of friction.

(i) State Newton's that law of motion which defines force as the agent of motion change.

(j) What is the graphical representation of the moment of a force ?

2. Answer the following questions :  $2 \times 5 = 10$

(a) Three equal forces acting at a point are in equilibrium. Show that they are equally inclined to one another.

(b) Find the position of centre of gravity (C.G) of a uniform semicircular arc of radius  $a$ .

(c) Prove that earth's gravitational field is a conservative force field.

(d) Two men have to carry a block of stone of weight  $70\text{kg}$  on a light plank. How must the block be placed so that one of the men should bear a weight of  $10\text{kg}$  more than the other ?

(e) Prove that the change in kinetic energy of a body is equal to the work done by the acting force.

3. Answer the following questions : **(any four)**

$5 \times 4 = 20$

(a) Two forces  $P$  and  $Q$  acting on a particle at an angle  $\alpha$  have a resultant

$(2k+1)\sqrt{P^2+Q^2}$ . When they act at an angle  $90^\circ - \alpha$ , the resultant becomes

$(2k-1)\sqrt{P^2+Q^2}$ , prove that

$$\tan \alpha = \frac{k-1}{k+1}.$$

(b) If the two like parallel forces  $P$  and  $Q$  acting on a rigid body at  $A$  and  $B$  be interchanged in position, then show that the point of application of the resultant will be displaced along  $\overline{AB}$  through a distance  $d$  where

$$d = \frac{P-Q}{P+Q} \cdot AB \quad (P > Q).$$

(c) Forces of magnitudes 1, 2, 3, 4,  $2\sqrt{2}$  act respectively along the sides  $\overline{AB}$ ,  $\overline{BC}$ ,  $\overline{CD}$ ,  $\overline{DA}$  and the diagonal  $\overline{AC}$  of the square  $ABCD$ . Show that their resultant is a couple, and find its moment.

- (d) A particle moves towards a centre of attraction starting from rest at a distance  $a$  from the centre. If its velocity when at any distance  $x$  from the centre

vary as  $\sqrt{\frac{a^2 - x^2}{x^2}}$ , find the law of force.

- (e) An elastic string without weight, of which the unstretched length is  $l$  and the modulus of elasticity is the weight of  $n$  ozs, is suspended by one end, and a mass of  $m$  ozs. is attached to the other; show that the time of a vertical

oscillation is  $2\pi\sqrt{\frac{ml}{ng}}$ .

- (f) A particle of mass  $m$  is projected vertically under gravity, the resistance of the air being  $mk$  times the velocity. Show that the greatest height attained

by the particle is  $\frac{V^2}{g}[\lambda - \log(1 + \lambda)]$ ,

where  $V$  is the terminal velocity of the particle and  $\lambda V$  is the initial vertical velocity.

4. Answer the following questions : **(any four)**

10×4=40

- (a) (i) Forces  $P, Q, R$  acting along  $\overline{IA}, \overline{IB}, \overline{IC}$ , where  $I$  is the in-centre of the triangle  $ABC$ , are in equilibrium. Show that 4

$$P : Q : R = \cos \frac{1}{2}A : \cos \frac{1}{2}B : \cos \frac{1}{2}C$$

- (ii) Forces  $L, M, N$  act along the sides of the triangle formed by the lines  $x + y - 1 = 0, x - y + 1 = 0, y = 2$ . Find the magnitude and the line of action of the resultant. 6

- (b) A body is resting on a rough inclined plane of inclination  $\alpha$  to the horizon, the angle of friction being  $\lambda (\alpha > \lambda)$ . If the body is acted on by a force  $P$ , then find the magnitude of  $P$  when

- (i) the body is just on the point of slipping down.

- (ii) the body is just on the point of sliding up.

- (c) (i) Find the C.G of the area of the cardioid  $r = a(1 + \cos \theta)$  5

(ii) Find the C.G. of the solid formed by the revolution of the quadrant of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  about its minor axis. 5

(d) (i) Three forces  $P, Q, R$  act in the same sense along the sides  $\overrightarrow{BC}, \overrightarrow{CA}, \overrightarrow{AB}$  of a triangle  $ABC$ . Show that, if their resultant passes through the centroid, then

$$P \operatorname{Cosec} A + Q \operatorname{Cosec} B + R \operatorname{Cosec} C = 0 \quad 5$$

(ii) Forces  $P, Q, R, S$  act along the sides  $\overrightarrow{AB}, \overrightarrow{BC}, \overrightarrow{CD}, \overrightarrow{DA}$  of the cyclic quadrilateral  $ABCD$ , taken in order, where  $A$  and  $B$  are the extremities of a diameter. If they are in equilibrium, then prove that

$$R^2 = P^2 + Q^2 + S^2 + \frac{2PQS}{R} \quad 5$$

(e) The velocities of a particle along and perpendicular to the radius from a fixed origin are  $\lambda r$  and  $\mu \theta$ . Find the path. Also show that the accelerations along and perpendicular to the radius vector

$$\text{are } \lambda^2 r - \frac{\mu^2 \theta^2}{r} \text{ and } \mu \theta \left( \lambda + \frac{\mu}{r} \right).$$

(f) A particle moves in a straight line  $OA$  with an acceleration which is always directed towards  $O$  and varies inversely as the square of its distance from  $O$ . If initially the particle were at rest at  $A$ , show that the time taken by it to arrive

$$\text{at the origin is } \frac{\pi a^{3/2}}{2\sqrt{2\mu}}.$$

(g) Show that the accelerations along the tangent and the normal to the path of

$$\text{a particle are } \frac{d^2 s}{dt^2} \left( = v \frac{dv}{ds} \right) \text{ and } \frac{v^2}{\rho},$$

where  $\rho$  is the radius of curvature of the curve at the point considered.

(h) A particle falls under gravity, supposed constant, in a resisting medium whose resistance varies as the square of the velocity. Discuss the motion, if the particle starts from rest.



### OPTION-C

Paper : MAT-HE-5036

#### (Probability and Statistics)

1. Answer the following questions :  $1 \times 10 = 10$

(a) If  $A$  and  $B$  are mutually exclusive then find  $P(A \cap B)$  and  $P(A \cup B)$ .

(b) Define probability mass function for discrete random variable.

(c) If  $P(x) = \frac{x}{15}$ ,  $x = 1$

0, elsewhere

Find  $P\{x = 1 \text{ or } x = 2\}$

(d) If  $X_1$  and  $X_2$  are independent random variables then what will be the modified statement of

$$V(X_1 + X_2) = V(X_1) + V(X_2) + 2\text{cov}(X_1, X_2)$$

(e) If a non-negative real valued function  $f$  is the probability density function of some continuous random variable, then

what is the value of  $\int_{-\alpha}^{\alpha} f(x) dx$  ?

(f) Name the discrete distribution for which mean and variance have the same value. What is the value ?

(g) What is meant by mathematical expectation of a random variable ?

(h) Under what condition the binomial distribution becomes the normal distribution.

(i) Write the equation of line of regression of  $y$  on  $x$ .

(j) State weak law of large number.

2. Answer the following questions :  $2 \times 5 = 10$

(a) If the events  $A$  and  $B$  are independent of  $A$  and  $B$  separately, is it necessary that they are independent of  $A \cap B$  ? Justify.

(b) Let  $X$  be a random variable with the following probability distribution :

$x:$	-3	6	3
$P(X = x):$	$\frac{1}{6}$	$\frac{1}{2}$	$\frac{1}{3}$

Find  $E(X^2)$

(c) State two properties of Poisson distribution.

- (d) If  $X$  is a continuous random variable whose probability density function is given by

$$f(x) = c(4x - 2x^2), 0 < x < 2 \\ = 0, \text{ otherwise}$$

then find the value of  $c$ .

- (e) If  $X$  is a random variable, then prove that  $\text{Var}(ax + b) = a^2 \text{Var}(X)$  where  $a$  and  $b$  are constants.

3. Answer **any four** parts from the following :

$$5 \times 4 = 20$$

- (a) A bag contains 6 white and 9 black balls. Four balls are drawn at a time. Find the probability for the first draw to give 4 white and the second to give 4 black balls if the balls are not replaced before the second draw.
- (b) The probability density function of a two dimensional random variable  $(X, Y)$  is given by

$$f(x, y) = x + y, 0 < x + y < 1 \\ 0, \text{ elsewhere}$$

Evaluate  $P\left(X < \frac{1}{2}, Y > \frac{1}{4}\right)$

- (c) A die is tossed twice. Getting 'a number greater than 4' is considered a success. Find the mean and variance of the probability distribution of the number of success.

- (d) The joint density function of two random variables  $X$  and  $Y$  is given by

$$f(x, y) = \frac{xy}{96}, 0 < x < 4, 1 < y < 5 \\ 0, \text{ otherwise}$$

Find

- (i)  $E(X)$
- (ii)  $E(Y)$
- (iii)  $E(2X + 3Y)$
- (e) For any two independent random variable  $X$  and  $Y$ , for which  $E(X)$  and  $E(Y)$  exists, show that
- $$E(XY) = E(X)E(Y)$$
- (f) With usual notation for a binomial variate  $X$ , given that  $9p(X=4) = p(X=2)$  when  $n = 6$   
Find the value of  $p$  and  $q$ .

4. Answer **any four** parts from the following :

$$10 \times 4 = 40$$

(a) (i) If  $A$  and  $B$  are any two events and are not disjoint then show that

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

Hence find  $P(A \cup B \cup C)$ .

(ii) From a bag containing 4 white and 6 red balls, three balls are drawn at random. Find the expected number of white balls drawn.

(b) (i) The joint density function of  $x$  and  $y$  is given by

$$f(x, y) = \begin{cases} 2e^{-x}e^{-2y}, & 0 < x < a, 0 < y < a \\ 0, & \text{otherwise} \end{cases}$$

compute  $P(X > 1, Y < 1)$ ,  $P(X < Y)$  and  $P(X < a)$ .

(ii) If  $X$  is a random Poisson variate with parameter  $m$ , then show that

$$p(X \geq n) - p(X \geq n+1) = \frac{e^{-m}m^n}{L^n}$$

(c) (i) If  $X$  is a binomial variate then prove that

$$\text{Cov}\left(\frac{X}{n}, \frac{n-X}{n}\right) = -\frac{pq}{n}$$

(ii) Show that normal distribution may be regarded as a limiting case of Poisson's distribution on the parameter  $m \rightarrow \infty$ .

(d) (i) Prove that the moment generating function of the sum of a number of independent random variables is equal to the product of their respective moment generating function.

(ii) Define moments and moment generating function of a random variable  $X$ . If  $M(t)$  is the moment generating function of a random variable  $X$  about the origin, show that the moment  $\mu'_r$  is given by

$$\mu'_r = \left[ \frac{d^r M(t)}{dt^r} \right]_{t=0}$$

(e) (i) If  $U = \frac{X-a}{h}$ ,  $V = \frac{Y-b}{k}$  where  $a, b, h, k$  are constants,  $h > 0, k > 0$  then show that  $r(X, Y) = r(U, V)$ .

( $r$  represents the correlation co-efficient)

- (ii) The two regression equations of the variables  $x$  and  $y$  are

$$x = 19.13 - 0.87y$$

$$y = 11.64 - 0.50x$$

Find (l) mean of  $x$ 's

(m) mean of  $y$ 's

(n) correlation co-efficients between  $x$  and  $y$ .

- (f) (i) Find the mean and variance of a Binomial distribution.

- (ii) If  $X$  is a random variable with mean  $\mu$  and variance  $\sigma^2$  then for any positive number  $k$ , prove that

$$P\{|X - \mu| \geq k\sigma\} \leq \frac{1}{k^2}$$

- (g) (i) A function  $f(x)$  of  $x$  is defined as follows :

$$\begin{aligned} f(x) &= 0 && \text{for } x < 2 \\ &= \frac{1}{18}(3 + 2x) && \text{for } 2 \leq x \leq 4 \\ &= 0 && \text{for } x > 4 \end{aligned}$$

Show that it is a density function. Also find the probability that a variate with this density will lie in the interval  $2 \leq x \leq 3$ .

- (ii) Two random variables  $X$  and  $Y$  have the following joint probability distribution function.

$$f(x, y) = \begin{cases} 2 - x - y, & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Find (l) marginal density function

(m)  $E(X)$  and  $E(Y)$

(n) conditional density function

- (h) (i) Show that Poisson distribution is a limiting case of the Negative Binomial Distribution.

- (ii) Let the random variable  $X_i$  assume values  $i$  and  $-i$  with equal probabilities. Show that the law of large number cannot be applied to the independent variables  $X_1, X_2, X_3, \dots, X_n$ .