

Total number of printed pages-11

3 (Sem-1/CBCS) MAT HC 1

2022

**MATHEMATICS**

(Honours)

Paper : MAT-HC-1016

(Calculus)

Full Marks : 60

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer **any seven** of the following questions : 1×7=7
- (a) Write down the  $n^{\text{th}}$  derivative of  $\cos(5x+3)$ .
- (b) Write when the graph of a function  $f$  is said to have vertical tangent at a point  $P(c, f(c))$ .

Contd.

(c) Write down the value of  $\lim_{x \rightarrow +\infty} x^n e^{-kx}$

(d) Evaluate  $\int_0^{\pi/2} \sin^6 x dx$

(e) In terms of marginal revenue and marginal cost, when is the profit maximized?

(f) For what purpose the disk and washer methods are used?

(g) Parameterize the curve  $y = 4x^2$

(h) When the graph of a vector function  $\vec{F}(t)$  is said to be smooth?

(i) Determine the values of  $t$  for which the vector function  $\vec{F}(t) = \frac{\hat{i} + 2\hat{j}}{t^2 + 1}$  is continuous?

(j) State the geometrical significance of the scalar triple product of vectors  $\vec{u}$ ,  $\vec{v}$  and  $\vec{w}$ .

(k) Find  $\int_0^{\pi} (t\hat{i} + 3\hat{j} - \sin t \hat{k}) dt$

(l) When a function  $f$  is said to be continuously differentiable on an interval  $I$ ?

2. Answer **any four** of the following questions :  
2×4=8

(a) Evaluate  $\lim_{x \rightarrow +\infty} \frac{3x^3 - 5x + 9}{5x^3 + 2x^2 - 7}$

(b) Using Leibnitz's rule obtain the  $n^{\text{th}}$  derivative of  $y = x^3 e^x$ .

(c) By integration find the length of the circle  $r = 2 \sin \theta$ .

(d) Let  $\vec{F}(t) = \hat{i} + e^t \hat{j} + t^2 \hat{k}$  and

$$\vec{G}(t) = 3t^2 \hat{i} + e^{-t} \hat{j} + 2t \hat{k},$$

then find  $\frac{d}{dt} \{ \vec{F}(t) \cdot \vec{G}(t) \}$ .

(e) Find the tangent vector to the graph of the vector function  $\vec{F}(t) = t^2 \hat{i} + 2t \hat{j} + e^t \hat{k}$  at the point  $t = -1$ .

(f) State Kepler's laws of motion.

(g) Find the volume generated by revolving about OX, the area bounded by  $y = x^3$  between  $x = 0$  and  $x = 2$ .

(h) Find the length of the polar curve  $r = e^{3\theta}$ ,  $0 \leq \theta \leq \pi/2$ .

3. Answer **any three** of the following :

$$5 \times 3 = 15$$

(a) Find the constants  $a$  and  $b$  that guarantee that the graph of the function

$$\text{defined by } f(x) = \frac{ax+5}{3-bx}.$$

will have a vertical asymptote at  $x = 5$  and a horizontal asymptote at  $y = 3$ .

(b) Evaluate :

$$2+3=5$$

(i)  $\lim_{x \rightarrow \pi/2^-} (x - \pi/2) \tan x$

(ii)  $\lim_{x \rightarrow +\infty} \left( 1 + \frac{1}{2x} \right)^{3x}$

(c) If  $I_n = \int_0^{\pi/4} \tan^n \theta d\theta$ , then prove that

$$n(I_{n+1} + I_{n-1}) = 1.$$

Hence evaluate  $\int_0^{\pi/4} \tan^3 \theta d\theta$ .  $3+2=5$

(d) A firm determines that  $x$  units of its product can be sold daily at  $p$  rupees per unit where  $x = 1000 - p$ . The cost of producing  $x$  units per day is  $C(x) = 3000 + 20x$ .

Find the revenue function  $R(x)$ .

Find the profit function  $P(x)$ .

Assuming that the production capacity is at most 500 units per day, determine how many units the company must produce and sell each day to maximize the profit.  $1+1+3=5$

- (e) Show that a cone of radius  $r$  and height  $h$  has lateral surface area

$$S = \pi r \sqrt{r^2 + h^2}.$$

- (f) For any three vectors  $\vec{a}, \vec{b}, \vec{c}$  in space, prove that  $\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{c} \cdot \vec{a}) \vec{b} - (\vec{b} \cdot \vec{a}) \vec{c}$ .

- (g) Use cylindrical shell method to find the volume of the solid generated when the region  $R$  under  $y^2 = x$  and  $x$ -axis over the interval  $[0, 4]$  is revolved about the line  $y = -1$ .

- (h) If the non-zero vector function  $\vec{F}(t)$  is differentiable and has constant length, then prove that  $\vec{F}(t)$  is orthogonal to the derivative vector  $\vec{F}'(t)$ .

Verify this result for

$$\vec{F}(t) = \cos t \hat{i} + \sin t \hat{j} + 3\hat{k}. \quad 3+2=5$$

4. Answer **any three** of the following :

$$10 \times 3 = 30$$

- (a) State Leibnitz's theorem. Use it to show that if  $y = e^{m \cos^{-1} x}$ , then

$$(1 - x^2) y_{n+2} - (2n+1) x y_{n+1} - (n^2 + m^2) y_n = 0$$

Hence find  $y_n(0)$ .  $2+5+3=10$

- (b) Find the vertical and horizontal asymptotes (if any) of the graph of the function

$$f(x) = \frac{x^2 - x - 2}{x - 3}.$$

Find where the graph is rising, where it is falling, determine concavity, locate all critical points and points of inflection. Finally sketch the graph.

- (c) Obtain the reduction formula for  $\int \sin^n x \, dx$ .

Hence evaluate

(i)  $\int_0^{\pi/2} \sin^n x \, dx$

(ii)  $\int_0^{\pi/2} \sin^7 x \, dx$

$5+3+2=10$

- (d) A boy standing at the edge of a cliff throws a ball upward at an angle of  $30^\circ$  with an initial speed of  $64 \text{ ft/s}$ . Suppose that when the ball leaves the boy's hand, it is  $48 \text{ ft}$  above the ground at the base of the cliff.

- (i) What are the time of flight of the ball and its range?  
 (ii) What are the velocity of the ball and its speed at impact?

- (iii) What is the highest point reached by the ball during its flight?

$3+3+4=10$

- (e) (i) Find the area of the surface generated by revolving about the  $x$ -axis the top half of the cardioid  $r = 1 + \cos \theta$ .

5

- (ii) Using disk method find the volume generated when the region bounded by the line  $y = 4 - x$  and the  $x$ -axis on the interval  $0 \leq x \leq 4$  revolve about the line  $x = -2$ .

5

- (f) (i) Find the position vector  $\vec{R}(t)$  and velocity vector  $\vec{V}(t)$ , given the acceleration  $\vec{A}(t)$  and initial position and velocity vectors  $\vec{R}(0)$  and  $\vec{V}(0)$  as

$\vec{A}(t) = t^2 \hat{i} - 2\sqrt{t} \hat{j} + e^{3t} \hat{k}$

$\vec{R}(0) = 2\hat{i} + \hat{j} - \hat{k}, \vec{V}(0) = \hat{i} - \hat{j} - 2\hat{k}$ .

5

(ii) A particle moves along the parametric curve  $x = 2t$ ,  $y = t$ . Find the position vector  $\vec{R}(t)$  and velocity vector  $\vec{V}(t)$  in terms of  $\hat{U}_r$  and  $\hat{U}_\theta$ . 5

(g) (i) It is projected that  $t$  years from now, the population of a certain country will be  $P(t) = 50e^{0.02t}$  million.

At what rate will the population be changing with respect to time 10 years from now?

At what percentage of rate, will the population be changing with respect to time  $t$  years from now?

3+3=6

(ii) Find the length of the curve defined by  $9x^2 = 4y^3$  between the points  $(0, 0)$  and  $(2\sqrt{3}, 3)$ . 4

(h) A object moving along a smooth curve has velocity  $\vec{v}$  given by  $\vec{v} = \frac{ds}{dt} \hat{T}$ .

Deduce the expression for acceleration

in the form  $\vec{A} = \frac{d^2s}{dt^2} \hat{T} + k \left( \frac{ds}{dt} \right)^2 \hat{N}$

where  $s$  is the arc length along the trajectory and  $k$  is the curvature. For an object moving along a helix with position vector  $\vec{R}(t) = (\cos t, \sin t, t)$  at any instant  $t$ , find the tangential and normal components of acceleration.

5+5=10