Total number of printed pages-7

3 (Sem-6/CBCS) MAT HC2

2023

MATHEMATICS

(Honours Core)

Paper: MAT-HC-6026

(Partial Differential Equation)

Full Marks: 60

Time: Three hours

The figures in the margin indicate full marks for the questions.

1. Answer the following:

 $1 \times 7 = 7$

- (i) The first order, quasi linear and linear partial differential equation are solved by using
 - (a) Lagrange's method
 - (b) Charpit's method

- (c) Jacobi method
- (d) None of the above (Choose the correct answer)
- (ii) The partial differential equation

$$x\left(\frac{\partial^2 z}{\partial x^2}\right) + \frac{\partial^2 z}{\partial y^2} = x^2$$
 is classified as

- (a) Parabolic, x = 0
- (b) Elliptic, x > 0
- (c) Hyperbolic, x < 0
- (d) All of the above (Choose the correct answer)
- (iii) What are the order and degree of $\frac{\partial^2 z}{\partial x^2} = \sqrt{1 + \frac{\partial z}{\partial y}}$?
- (iv) What type of partial differential equation is readily solved by Charpit's method?

- (v) The equation $p^2 + q^2 = 1$ is
 - (a) linear
 - (b) semi linear
 - (c) quasi linear
 - (d) Non-linear
 (Choose the correct answer)
- (vi) The solution which has number of arbitrary constants equal to number of independent variables is
 - (a) general integral
 - (b) complete integral
 - (c) particular integral
 - (d) singular integral
 (Choose the correct answer)
- (vii) Write down the form obtained of the PDE, in a function X(x, y) and two variables x, y after separation of variables is applied.

3

2. Answer in short: 2×4=8

- (i) Write down the construction of a first order partial differential equation.
- (ii) Define partial differential equation. Give one example.
- (iii) Eliminate arbitrary constants from $z = Ae^{pt} \sin px$ to form a partial differential equation.
- (iv) Determine whether the given equation is parabolic, elliptic or hyperbolic

$$y^2 \frac{\partial^2 z}{\partial x^2} - x^2 \frac{\partial^2 z}{\partial y^2} = 0$$

3. Answer any three: 5×3=15

(i) Eliminate the arbitrary function f from the equation $f(x^2 + y^2 + z^2, z^2 - xy) = 0$

- (ii) Find the general integrals of the linear partial differential equations $z(xp-yq)=y^2-x^2$
- (iii) Find the equation of the integral surface of the differential equation 2y(z-3)p + (2x-z)q = y(2x-3) which passes through the circle z = 0, $x^2 + y^2 = 2x.$
- (iv) Reduce to canonical form and find the general solution of $u_x + u_y = u$.
- (v) Apply the method of separation of variables u(x, y) = f(x) g(y) to solve the equation $y^2 u_x^2 + x^2 u_y^2 = (xyu)^2$.
- 4. Answer the following questions: 10×3=30
 - (a) Find a complete integral of $(p^2 + q^2)y = qz$ by Charpit's method.

5

Or

Apply the method of seperation of variables u(x, y) = f(x)g(y) to solve the equation $u_x + 2u_y = 0$, $u(0, y) = 3e^{-2y}$.

(b) Solve $p_3x_3(p_1 + p_2) + x_1 + x_2 = 0$ by Jacobi method.

The wat of nontrine language

Transform the equation to canonical form $u_{xx} + y^2 u_{yy} = y$.

(c) Obtain the general solution of the equation

$$x^{2}u_{xx} + 2xyu_{xy} + y^{2}u_{yy} + xyu_{x} + y^{2}u_{y} = 0$$

Solve the following:

(i)
$$x(y^2-z^2)p+y(z^2-x^2)q=z(x^2-y^2)$$

(ii)
$$(x^2 - y^2 - z^2)p + 2xyq = 2xz$$

7