

Total number of printed pages-7

3 (Sem-4/CBCS) MAT HC 2

2024

**MATHEMATICS**

(Honours Core)

Paper : MAT-HC-4026

**( Numerical Methods )**

Full Marks : 60

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer the following as directed : 1×7=7
- (a) Name the three basic components of an algorithm.
  - (b) Show  $\nabla E \equiv \Delta$ .
  - (c) Write down the Lagrangian linear interpolation formula at the points  $x_0$  and  $x_1$  with corresponding function values  $f_0$  and  $f_1$ .

Contd.

- (d) What is the order of convergence of secant method?
- (e) The approximation formula for finding the derivative at  $x_0$  given by

$$f'(x_0) = \frac{f(x_0 + h) - f(x_0)}{h} - \frac{h}{2} f''(\xi),$$

$$x_0 < \xi < x_0 + h$$

is a

- (i) backward difference approximation formula of first order of approximation
- (ii) forward difference approximation formula of second order of approximation
- (iii) forward difference approximation formula of first order of approximation
- (iv) None of the above  
(Choose the correct option)
- (f) What is numerical integration? What is its general form?
- (g) Name a method for approximating a solution to an initial value problem.

2. Answer the following questions :  $2 \times 4 = 8$

- (a) Compute the following limit and determine the rate of convergence

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{x}$$

- (b) Prove  $(I + \Delta)(I - \nabla) \equiv I$ .

- (c) Show that LU decomposition of a matrix is unique up to scaling by a diagonal matrix.

- (d) Find the approximate value of  $\int_0^1 \frac{dx}{1+x}$  by Simpson's rule.

3. Answer **any three** :  $5 \times 3 = 15$

- (a) Construct an iteration function corresponding to the given function

$$f(x) = x^3 - x^2 - 10x + 7.$$

Use the fixed point iteration scheme with initial approximation as  $P_0 = 1$  and perform three iterations to approximate the root of  $f(x) = 0$ .

- (b) Using the data given below form the divided difference table and use it to construct the Newton form of the interpolating polynomial :

$x$	-1	0	1	2
$y$	5	1	1	11

- (c) Use four iterations of Newton's method to approximate the root of the equation

$$f(x) = x^3 + 2x^2 - 3x - 1$$

in the interval (1, 2) starting with an initial approximation of  $P_0 = 1$ .

- (d) Derive the second order central difference approximation for first derivative including error term given by

$$f'(x_0) = \frac{f(x_0 + h) - f(x_0 - h)}{2h} - \frac{h^2}{6} f'''(\xi)$$

- (e) (i) Name the measures by which errors are quantified. Write down the expressions for the same.

(ii) Prove that  $\Delta^n f(x_i) = (E - I)^n f(x_i)$

4. Answer **any three** : 10×3=30

- (a) What is Theoretical Error Bound ? Show that the Bisection Method for approximating a root of the equation  $f(x) = 0$  always converges. Find the order of convergence of the Bisection Method. 1+6+3=10

- (b) Verify that the equation  $x^3 + x^2 - 3x - 3 = 0$  has a root in the interval (1, 2). Given that the exact root is  $x = \sqrt{3}$ , perform the first three iterations of the Regula-Falsi method. What is the computable estimate for  $|e_n|$ , the error obtained in  $n$ th step by this method. Verify that the absolute error in the third approximation satisfies the error estimate. 1+6+3=10

- (c) What is an interpolating polynomial? Determine the interpolation error when a function is approximated by a constant polynomial. Mention an advantage and a disadvantage of Lagrangian form of the interpolating polynomial. Derive the Lagrangian interpolating polynomial for the given data : 1+2+2+5=10

$x$	-2	-1	0	1	2	3
$y$	39	3	-1	-3	-9	-1

- (d) What are two different classes of methods for solving a linear system of equations. Name one method of each type. What do you mean by an LU decomposition of square matrix A.

Solve the following system using LU decomposition :  $1+1+8=10$

$$2x_1 + 7x_2 + 5x_3 = -4$$

$$6x_1 + 20x_2 + 10x_3 = -16$$

$$4x_1 + 3x_2 = -7$$

- (e) (i) Derive the basic Trapezoidal rule

for integrating  $\int_a^b f(x) dx$ . 6

- (ii) Use appropriate first order approximation formulas to find derivatives of the values of  $f(x)$  at the points  $x = 0.5$ ,  $x = 0.6$  and  $x = 0.7$ . 4

$x$	$f(x)$	$f'(x)$
0.5	0.4794	?
0.6	0.5646	?
0.7	0.6442	?

- (f) What is the basic problem that is solved by Euler's method? Derive Euler's method. Given that the exact solution

to  $\frac{dx}{dt} = \frac{t}{x}$  is  $x(t) = \sqrt{t^2 + 1}$ , find the

absolute error at each step that is obtained by solving

$$\frac{dx}{dt} = \frac{t}{x}, \quad 0 \leq t \leq 1.0, \quad x(0) = 1, \quad h = 0.5$$

by Euler's method. 1+4+5=10

---